

BEST AVAILABLE COPY

SIXTH EDITION

Organic Chemistry

Robert Thornton Morrison

Robert Neilson Boyd

New York University




Prentice Hall, Englewood Cliffs, New Jersey 07632


[illegible]

20.2 Nomenclature


The names of acid derivatives are taken in simple ways from either the common name or the IUPAC name of the corresponding carboxylic acid. For example:




Acetic acid
Ethanoic acid



Benzoic acid




Acetyl chloride
Ethanoyl chloride




Benzoyl chloride

Change:
-ic acid to -yl chloride




Acetic anhydride
Ethanoic anhydride




Benzoic anhydride

acid to anhydride




Acetamide
Ethanamide




Benzamide

-ic acid of common name
(or -oic acid of IUPAC name)
to -amide



Ethyl acetate
Ethyl ethanoate




Ethyl benzoate

-ic acid to -ate,
preceded by name of
alcohol or phenol group

20.3 Physical properties

The presence of the C=O group makes the acid derivatives polar compounds. Acid chlorides and anhydrides (Table 20.1) and esters (Table 20.2, p. 769) have boiling points that are about the same as those of aldehydes or ketones of comparable molecular weight (see Sec. 18.3). Amides (Table 20.1) have quite high boiling points because they are capable of strong intermolecular hydrogen bonding.



The b
the esters t
the usual o
Volati
in the prep
irritating o
acids.

Acetyl cl
Propiony
n-Butyryl
n-Valeryl
Stearoyl cl
Benzoyl cl
p-Nitrober
chloride
3,5-Dinitro
chloride

Acetic anh
Phthalic ar

Before v
outline certa
individual fa
Each de
correspondin
acid by simpl
conversion o
has certain cl

The deri
carbonyl gro
undergone by
But by its pre
these compou

Here, to
functions: (a)
acidity of hyd
(We shal
21.11-21.12 a